

BMP4

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:100 - 1:200

Calculated MW:47kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human BMP4

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

prt

Synonym:

BMP2B; BMP2B1; MCOPS6; ZYME;

Catalog #:A1565

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:652

Isotype:IgG

Swiss Prot:P12644

Purity:Affinity purification

For research use only.

Background:

Bone morphogenetic proteins (BMPs) were first identified as molecules that can induce ectopic bone and cartilage formation (1,2). BMPs belongs to the TGF- superfamily, playing many diverse functions during development (3). BMPs are synthesized as precursor proteins and then processed by cleavage to release the c-terminal mature BMP. BMPs initiate signaling by binding to a receptor complex containing type I and type II serine/threonine receptor kinases that then phosphorylate Smad (mainly Smad1, 5 and 8), resulting the translocation of Smad into the nucleus. BMP was also reported to activate MAPK pathways in some systems (3,4).BMP4 plays an essential role during development. Mouse embryos deficient in BMP4 die around the time of gastrulation (5). BMP4 regulates stem cell maintenance and differentiation in various systems (6-8).

To place an order, please [Click HERE](#).