

KIR3DL1

Reactivity:Human Mouse

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:49kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human KIR3DL1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

KIR; NKB1; NKAT3; NKB1B; CD158E1; KIR3DL1/S1;

Catalog #:A1617

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:3811

Isotype:IgG

Swiss Prot:P43629

Purity:Affinity purification

For research use only.

Background:

Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response.

To place an order, please [Click HERE](#).