

CTNND1

Reactivity:Human Mouse Rat

Tested applications:WB IHC ICC IF IP

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IF 1:50 - 1:200
IP 1:50 - 1:100

Calculated MW:105kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human CTNND1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

t

Synonym:

CAS; CTNND; KIAA0384; P120CAS; P120CTN; p120;

Catalog #:A1641

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:1500

Isotype:IgG

Swiss Prot:O60716

Purity:Affinity purification

For research use only.

Background:

Catenin -1 (p120 catenin) has an amino-terminal coiled-coil domain followed by a regulatory domain containing multiple phosphorylation sites and a central Armadillo repeat domain of ten linked 42-amino acid repeats. The carboxy-terminal tail has no known function (1). Catenin -1 fulfills critical roles in the regulation of cell-cell adhesion as it regulates E-cadherin turnover at the cell surface to determine the level of E-cadherin available for cell-cell adhesion (2). Catenin -1 has both positive and negative effects on cadherin-mediated adhesion (3). Actin dynamics are also regulated by catenin -1, which modulates RhoA, Rac, and cdc42 proteins (1). Analogous to -catenin, catenin -1 translocates to the nucleus, although its role at this location is unclear. Many studies show that catenin -1 is expressed irregularly or is absent in various types of tumor cells, suggesting that catenin -1 may function as a tumor suppressor (4). Phosphorylation of Tyr904 on Catenin--1 was identified at Cell Signaling Technology (CST) using PhosphoScan®, a CST's LC-MS/MS platform for phosphorylation site discovery (5).

To place an order, please [Click HERE](#).