

BARD1

Reactivity:Human Mouse Rat

Tested applications:WB

Recommended Dilution:WB 1:500 - 1:2000

Calculated MW:87kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human BARD1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Catalog #:A1685

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:580

Isotype:IgG

Swiss Prot:Q99728

Purity:Affinity purification

For research use only.

Background:

This gene encodes a protein which interacts with the N-terminal region of BRCA1. In addition to its ability to bind BRCA1 in vivo and in vitro, it shares homology with the 2 most conserved regions of BRCA1: the N-terminal RING motif and the C-terminal BRCT domain. The RING motif is a cysteine-rich sequence found in a variety of proteins that regulate cell growth, including the products of tumor suppressor genes and dominant protooncogenes. This protein also contains 3 tandem ankyrin repeats. The BARD1/BRCA1 interaction is disrupted by tumorigenic amino acid substitutions in BRCA1, implying that the formation of a stable complex between these proteins may be an essential aspect of BRCA1 tumor suppression. This protein may be the target of oncogenic mutations in breast or ovarian cancer. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2013]

To place an order, please [Click HERE](#).