www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## USF1

Reactivity: Human Mouse Rat

Tested applications: WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:34kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human USF1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

k

Synonym:

UEF; FCHL; MLTF; FCHL1; MLTFI; HYPLIP1; bHLHb11;

Background:

This gene encodes a member of the basic helix-loop-helix leucine zipper family, and can function as a cellular transcription factor. The encoded protein can activate transcription through pyrimidine-rich initiator (Inr) elements and E-box motifs. This gene has been linked to familial combined hyperlipidemia (FCHL). Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Jul 2008]

To place an order, please Click HERE.

Polyclonal Antibody

Species:Rabbit

Gene ID:7391

Isotype:IgG

Swiss Prot:P22415

Purity: Affinity purification

For research use only.



