

HLA-DPA1

Reactivity:Human

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:29kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human HLA-DPA1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

PLT1; HLADP; HLASB; DP(W3); DP(W4); HLA-DP1A;

Catalog #:A1754

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:3113

Isotype:IgG

Swiss Prot:P20036

Purity:Affinity purification

For research use only.

Background:

HLA-DPA1 belongs to the HLA class II alpha chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DPA) and a beta (DPB) chain, both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The alpha chain is approximately 33-35 kDa and its gene contains 5 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and the cytoplasmic tail. Within the DP molecule both the alpha chain and the beta chain contain the polymorphisms specifying the peptide binding specificities, resulting in up to 4 different molecules.

To place an order, please [Click HERE](#).