

HCFC1

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW: 209kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human HCFC1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

h

Synonym:

CFF; HCF1; HFC1; MRX3; VCAF; HCF-1;

Catalog #: A1841

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 3054

Isotype: IgG

Swiss Prot: P51610

Purity: Affinity purification

For research use only.

Background:

This gene is a member of the host cell factor family and encodes a protein with five Kelch repeats, a fibronectin-like motif, and six HCF repeats, each of which contains a highly specific cleavage signal. This nuclear coactivator is proteolytically cleaved at one of the six possible sites, resulting in the creation of an N-terminal chain and the corresponding C-terminal chain. The final form of this protein consists of noncovalently bound N- and C-terminal chains. The protein is involved in control of the cell cycle and transcriptional regulation during herpes simplex virus infection. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized. [provided by RefSeq, Jul 2008]

To place an order, please [Click HERE](#).