

## LIG3

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC IF

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:** 113kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human LIG3

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

e

**Synonym:**

LIG2; LIG3;

**Catalog #:** A1887

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 3980

**Isotype:** IgG

**Swiss Prot:** P49916

**Purity:** Affinity purification

For research use only.

**Background:**

This gene is a member of the DNA ligase family. Each member of this family encodes a protein that catalyzes the joining of DNA ends but they each have a distinct role in DNA metabolism. The protein encoded by this gene is involved in excision repair and is located in both the mitochondria and nucleus, with translation initiation from the upstream start codon allowing for transport to the mitochondria and translation initiation from a downstream start codon allowing for transport to the nucleus. Additionally, alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008]

**To place an order, please [Click HERE](#).**