

TNNC1

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:18kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human TNNC1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

CMD1Z; TNC; TNNC;

Catalog #:A1927

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:7134

Isotype:IgG

Swiss Prot:P63316

Purity:Affinity purification

For research use only.

Background:

Troponin, working in conjunction with tropomyosin, functions as a molecular switch, regulating muscle contraction in response to changes in the intracellular Ca²⁺ concentration. Troponin consists of three subunits: the Ca²⁺-binding subunit troponin C (TnC), the tropomyosin-binding subunit troponin T (TnT), and the inhibitory subunit troponin I (TnI) (1). In response to -adrenergic stimulation of the heart, Ser23 and Ser24 of TnI (cardiac) are phosphorylated by PKA and PKC. This phosphorylation stimulates a conformational change of the regulatory domain of TnC, reduces the association between TnI and TnC, and decreases myofilament Ca²⁺ sensitivity by reducing the Ca²⁺ binding affinity of TnC (1-3).

To place an order, please [Click HERE](#).