

## RPS27A

**Reactivity:**Human Mouse Rat

**Tested applications:**WB

**Recommended Dilution:**WB 1:500 - 1:2000

**Calculated MW:**9kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human RPS27A

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

j

**Synonym:**

CEP80; HUBCEP80; UBA80; UBCEP1; UBCEP80; UBC; S27A; RPS27A; Ubiquitin;

**Catalog #:**A2027

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**6233

**Isotype:**IgG

**Swiss Prot:**P62979

**Purity:**Affinity purification

For research use only.

**Background:**

Ubiquitin, a highly conserved protein that has a major role in targeting cellular proteins for degradation by the 26S proteasome, is synthesized as a precursor protein consisting of either polyubiquitin chains or a single ubiquitin fused to an unrelated protein. This gene encodes a fusion protein consisting of ubiquitin at the N terminus and ribosomal protein S27a at the C terminus. When expressed in yeast, the protein is post-translationally processed, generating free ubiquitin monomer and ribosomal protein S27a. Ribosomal protein S27a is a component of the 40S subunit of the ribosome and belongs to the S27AE family of ribosomal proteins. It contains C4-type zinc finger domains and is located in the cytoplasm. Pseudogenes derived from this gene are present in the genome. As with ribosomal protein S27a, ribosomal protein L40 is also synthesized as a fusion protein with ubiquitin; similarly, ribosomal protein S30 is synthesized as a fusion protein with the ubiquitin-like protein fubi. Multiple alternatively spliced transcript variants that encode the same proteins have been identified.[provided by RefSeq, Sep 2008]

**To place an order, please [Click HERE](#).**