

## SLAMF1

**Reactivity:**Human

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**70kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human SLAMF1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

CD150; CDw150; SLAM;

**Catalog #:**A2044

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**6504

**Isotype:**IgG

**Swiss Prot:**Q13291

**Purity:**Affinity purification

For research use only.

**Background:**

Following occupancy of the T cell receptor by antigen, T cell proliferation and lymphokine production are determined by a second costimulatory signal delivered by a ligand expressed on antigen-presenting cells. SLAM (for signaling lymphocyte-activation molecule, also designated CDw150) is a novel receptor on T cells that, when engaged, potentiates T cell expansion in a CD28-independent manner. SAP (for SLAM-associated protein) contains an SH2 domain and functions to inhibit SH-PTP2 recruitment to the SLAM docking site, an activity induced by Fyn phosphorylation of SLAM. Mutations of the SAP gene may be associated with X-linked lympho-proliferative disease (XLP).

**To place an order, please [Click HERE](#).**