

## IL13

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:100

**Calculated MW:**16kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human IL13

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

ALRH; BHR1; IL-13; MGC116786; MGC116788; MGC116789; P600;

**Background:**

IL-13 is produced by T cells and is important in the TH2 response. IL-13 targets include B cells, eosinophils, fibroblasts, mast cells and macrophages (1-3). IL-13 binds specifically to IL-13R1 that complexes with IL-4R to form the Type II IL-4R. Jak1 and Tyk2 are activated and signal through Stat3 and Stat6 (6). IL-13R2 is a different gene product, lacks the intracellular domain, does not complex with IL-4R and does not signal (1,6,7). The extracellular domain of IL-13R2 is often elevated in diseased states. IL-13 plays key roles in airway hyperresponsiveness (AHR) of allergic asthma (1,4,5) and modulates resistance to parasitic organisms (1).

**To place an order, please [Click HERE](#).**

**Catalog #:**A2089

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**3596

**Isotype:**IgG

**Swiss Prot:**P35225

**Purity:**Affinity purification

For research use only.