

ZYX

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW: 62kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human ZYX

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

j

Synonym:

Zyxin; ESP-2; HED-2;

Catalog #: A2135

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 7791

Isotype: IgG

Swiss Prot: Q15942

Purity: Affinity purification

For research use only.

Background:

The zyxin family of proteins includes LIMD1, ajuba, trip6 and zyxin, each of which contains three LIM domains at the carboxy-terminus. Zyxin family members associate with the actin cytoskeleton and are components of both the cell-cell junction adhesive complex and the integrin-mediated adhesive complex. They shuttle in and out of the nucleus where they may function in transcriptional activation (1). Zyxin is involved in the regulation of mechanical force-induced actin polymerization at focal adhesions (2), and in regulation of adhesion and migration, possibly through recruitment of Ena/VASP proteins to focal adhesions (3). Zyxin interacts with and may regulate the function of the tumor suppressor myopodin, which inhibits tumor growth and metastasis (4).

To place an order, please [Click HERE](#).