

IL10

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:100

Calculated MW: 19kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human IL10

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

b

Synonym:

IL10; CSIF; GVHDS; IL-10; IL10A; TGIF; Interleukin-10; Cytokine synthesis inhibitory factor;

Catalog #: A2171

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 3586

Isotype: IgG

Swiss Prot: P22301

Purity: Affinity purification

For research use only.

Background:

IL-10 is an anti-inflammatory cytokine that is produced by T cells, NK cells and macrophages (1,2). IL-10 initiates signal transduction by binding to a cell surface receptor complex consisting of IL-10 RI and IL-10 RII (1). Binding of IL-10 leads to the activation of Jak1 and Tyk2, which phosphorylates Stat-3 (1,3). The anti-inflammatory activity of IL-10 is due to its ability to block signaling through other cytokine receptors, notably IFN receptor, by upregulating expression of SOCS-1 (1,3). In addition, IL-10 promotes T cell tolerance by inhibiting tyrosine phosphorylation of CD28 (4,5). IL-10 is an important negative regulator of the immune response, which allows for maintenance of pregnancy (1). In contrast, increased IL-10 levels contribute to persistent Leishmania major infections (6).

To place an order, please [Click HERE](#).