

## IFIH1

**Reactivity:**Human Mouse

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:**117kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human IFIH1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

b

**Synonym:**

HLcd; IDDM19; MDA-5; MDA5; MGC133047;

**Catalog #:**A2203

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**64135

**Isotype:**IgG

**Swiss Prot:**Q9BYX4

**Purity:**Affinity purification

For research use only.

**Background:**

cts viral dsRNA and two amino-terminal caspase recruitment domains (CARD) that are required for triggering downstream signaling (4-7). Rig-I binds both dsRNA and viral ssRNA that contains a 5'-triphosphate end not seen in host RNA (8,9). Though structurally related, Rig-I and MDA-5 detect a distinct set of viruses (10,11). The CARD domain of the helicases, which is sufficient to generate signaling and IFN production, is recruited to the CARD domain of the MAVS/VISA/Cardif/IPS-1 mitochondrial protein, which triggers activation of NF-B, TBK1/IKK, and IRF-3/IRF-7 (12-15).

*To place an order, please [Click HERE](#).*