

5mC

Reactivity: Human Mouse Rat

Tested applications: ICC MeDIP

Recommended Dilution: ICC 1:50 - 1:200 MeDIP 1:1000 - 1:2000

Observed MW: Refer to Figures

Immunogen:

A synthetic peptide of human 5mC

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Catalog #: A2341

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Isotype: IgG

Purity: Affinity purification

For research use only.

Background:

5-Methylcytosine is the methylated form of the DNA base cytosine. In mammalian cells, DNA methylation is an enzymatic modification at the 5-position of cytosine present abundantly within the CpG dinucleotide sequence context. This DNA modification is inheritable and reversible without primary DNA base sequence changes, resulting in possible epigenetic modulation of phenotype and gene expression (1). CpG methylation may directly interfere with the binding of certain transcriptional regulators to their cognate DNA sequences or may enable recruitment of methyl-CpG binding proteins that create a repressed chromatin environment (2). The de novo formation and maintenance of 5-methylcytosine is catalyzed by DNA methyltransferase proteins (DNMTs) (1).

To place an order, please [Click HERE](#).