

MonoMethyl-Histone H4-K20

Reactivity:Human Mouse Rat Other (Wide Range)

Tested applications:WB IHC IF IP CHIP CHIPseq

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200 IP 1:50 - 1:200

ChIP 1:20 - 1:100 CHIPseq 1:20 - 1:100

Calculated MW:11kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human MonoMethyl-Histone H4-K20

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

u

Synonym:

H4K20me1; H4; H4/n; H4F2; H4FN; FO108; HIST2H4

Catalog #:A2370

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:8370

Isotype:IgG

Swiss Prot:P62805

Purity:Affinity purification

For research use only.

Background:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in a histone cluster on chromosome 1. This gene is one of four histone genes in the cluster that are duplicated; this record represents the centromeric copy.

To place an order, please [Click HERE](#).