

## PSMA1

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

**Calculated MW:**29kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human PSMA1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

NU; HC2; PROS30;

**Catalog #:**A2523

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**5682

**Isotype:**IgG

**Swiss Prot:**P25786

**Purity:**Affinity purification

For research use only.

**Background:**

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

**To place an order, please [Click HERE](#).**