

TNK2

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:114kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human TNK2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

ACK; ACK1; ACK-1; p21cdc42Hs;

Background:

This gene encodes a tyrosine kinase that binds Cdc42Hs in its GTP-bound form and inhibits both the intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs. This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3 domain. The protein may be involved in a regulatory mechanism that sustains the GTP-bound active form of Cdc42Hs and which is directly linked to a tyrosine phosphorylation signal transduction pathway. Several alternatively spliced transcript variants have been identified from this gene, but the full-length nature of only two transcript variants has been determined.

To place an order, please [Click HERE](#).

Catalog #:A2526

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:10188

Isotype:IgG

Swiss Prot:Q07912

Purity:Affinity purification

For research use only.