www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

AAK1

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:200 - 1:1000 IHC 1:10 - 1:50 Calculated MW:104kDa Observed MW:Refer to figures Immunogen: Recombinant protein of human AAK1 Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.



Adaptor-related protein complex 2 (AP-2 complexes) functions during receptor-mediated endocytosis to trigger clathrin assembly, interact with membrane-bound receptors, and recruit encodytic accessory factors. This gene encodes a member of the SNF1 subfamily of Ser/Thr protein kinases. The protein interacts with and phosphorylates a subunit of the AP-2 complex, which promotes binding of AP-2 to sorting signals found in membrane-bound receptors and subsequent receptor endocytosis. Its kinase activity is stimulated by clathrin. Alternatively spliced transcript variants have been described, but their biological validity has not been determined.

To place an order, please Click HERE.



Catalog #:A2652 Antibody Type: Polyclonal Antibody Species:Rabbit Gene ID:22848 Isotype:IgG Swiss Prot:Q2M2I8 Purity:Affinity purification

For research use only.



