

DAP3

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:200 - 1:1000 IHC 1:5 - 1:20

Calculated MW: 46kDa

Observed MW: Refer to figures

Immunogen:

Recombinant protein of human DAP3

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

bhj

Synonym:

DAP-3; MRPS29; MRP-S29; bMRP-10;

Catalog #: A2714

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 7818

Isotype: IgG

Swiss Prot: P51398

Purity: Affinity purification

For research use only.

Background:

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that also participates in apoptotic pathways which are initiated by tumor necrosis factor-alpha, Fas ligand, and gamma interferon. This protein potentially binds ATP/GTP and might be a functional partner of the mitoribosomal protein S27. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. Pseudogenes corresponding to this gene are found on chromosomes 1q and 2q.

To place an order, please [Click HERE](#).