www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## **FADD**

Reactivity: Human Mouse Rat

Tested applications:WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW:23kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human FADD

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

bis

Synonym:

GIG3; MORT1;

## Background:

The protein encoded by this gene is an adaptor molecule that interacts with various cell surface receptors and mediates cell apoptotic signals. Through its C-terminal death domain, this protein can be recruited by TNFRSF6/Fas-receptor, tumor necrosis factor receptor, TNFRSF25, and TNFSF10/TRAIL-receptor, and thus it participates in the death signaling initiated by these receptors. Interaction of this protein with the receptors unmasks the N-terminal effector domain of this protein, which allows it to recruit caspase-8, and thereby activate the cysteine protease cascade. Knockout studies in mice also suggest the importance of this protein in early T cell development.

To place an order, please Click HERE.

Polyclonal Antibody

Species: Rabbit

Gene ID:8772

Isotype:IgG

Swiss Prot:Q13158

Purity: Affinity purification

For research use only.





