

CCL13

Reactivity:Human

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100

Calculated MW:11kDa

Observed MW:Refer to figures

Immunogen:

A synthetic peptide of human CCL13

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

NCC1; CKb10; MCP-4; NCC-1; SCYL1; SCYA13;

Catalog #:A2853

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:6357

Isotype:IgG

Swiss Prot:Q99616

Purity:Affinity purification

For research use only.

Background:

This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils, but not neutrophils. This chemokine plays a role in accumulation of leukocytes during inflammation. It may also be involved in the recruitment of monocytes into the arterial wall during atherosclerosis.

To place an order, please [Click HERE](#).