

CCL17

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:11kD

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human CCL17

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

TARC; ABCD-2; SCYA17; A-152E5.3;

Catalog #:A2854

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:6361

Isotype:IgG

Swiss Prot:Q92583

Purity:Affinity purification

For research use only.

Background:

This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for T lymphocytes, but not monocytes or granulocytes. The product of this gene binds to chemokine receptors CCR4 and CCR8. This chemokine plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells.

To place an order, please [Click HERE](#).