www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

GAD1

Reactivity: Human Mouse Rat

Tested applications:WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW:67kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human GAD1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

GAD; SCP; CPSQ1;

Species: Rabbit

Gene ID:2571

Isotype:IgG

Swiss Prot:Q99259

Purity: Affinity purification

For research use only.

Background:

The enzyme glutamate decarboxylase (GAD) is responsible for the synthesis of the essential neurotransmitter gamma-aminobutyric acid (GABA) from L-glutamic acid (1). GAD1 (GAD67) and GAD2 (GAD65) are expressed in nervous and endocrine systems (2) and are thought to be involved in synaptic transmission (3) and insulin secretion (4), respectively. Autoantibodies against GAD2 may serve as markers for type I diabetes (5). Many individuals suffering from an adult onset disorder known as Stiff Person Syndrome (SPS) also express autoantibodies to GAD2 (6). Mutations in the GAD1 gene can cause autosomal recessive spastic cerebral palsy, possibly attributable to altered glutamate/GABA ratios (7).

To place an order, please Click HERE.





