

## KCNA1

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:100

**Calculated MW:** 56kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A synthetic peptide of human KCNA1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

g

**Synonym:**

EA1; MK1; AEMK; HBK1; HUK1; MBK1; RBK1; KV1.1

**Catalog #:** A2992

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 3736

**Isotype:** IgG

**Swiss Prot:** Q09470

**Purity:** Affinity purification

For research use only.

**Background:**

This gene encodes a voltage-gated delayed potassium channel that is phylogenetically related to the Drosophila Shaker channel. The encoded protein has six putative transmembrane segments (S1-S6), and the loop between S5 and S6 forms the pore and contains the conserved selectivity filter motif (GYGD). The functional channel is a homotetramer. The N-terminus of the channel is associated with beta subunits that can modify the inactivation properties of the channel as well as affect expression levels. The C-terminus of the channel is complexed to a PDZ domain protein that is responsible for channel targeting. Mutations in this gene have been associated with myokymia with periodic ataxia (AEMK).

**To place an order, please [Click HERE](#).**