

CCL8

Reactivity:Human

Tested applications:WB

Recommended Dilution:WB 1:200 - 1:1000

Calculated MW:11kDa

Observed MW:Refer to figures

Immunogen:

A synthetic peptide of human CCL8

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

HC14; MCP2; MCP-2; SCYA8; SCYA10;

Catalog #:A3008

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:6355

Isotype:IgG

Swiss Prot:P80075

Purity:Affinity purification

For research use only.

Background:

This antimicrobial gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils. By recruiting leukocytes to sites of inflammation this cytokine may contribute to tumor-associated leukocyte infiltration and to the antiviral state against HIV infection.

To place an order, please [Click HERE](#).