

SREBF1

Reactivity:Human

Tested applications:WB IHC FC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100 FC1:20 - 1:50

Calculated MW:125kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human SREBF1

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

k

Synonym:

SREBP1; bHLHd1; SREBP-1c;

Catalog #:A3139

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:6720

Isotype:IgG

Swiss Prot:P36956

Purity:Affinity purification

For research use only.

Background:

This gene encodes a transcription factor that binds to the sterol regulatory element-1 (SRE1), which is a decamer flanking the low density lipoprotein receptor gene and some genes involved in sterol biosynthesis. The protein is synthesized as a precursor that is attached to the nuclear membrane and endoplasmic reticulum. Following cleavage, the mature protein translocates to the nucleus and activates transcription by binding to the SRE1. Sterols inhibit the cleavage of the precursor, and the mature nuclear form is rapidly catabolized, thereby reducing transcription. The protein is a member of the basic helix-loop-helix-leucine zipper (bHLH-Zip) transcription factor family. This gene is located within the Smith-Magenis syndrome region on chromosome 17. Two transcript variants encoding different isoforms have been found for this gene.

To place an order, please [Click HERE](#).