

MNDA

Reactivity:Human

Tested applications:WB

Recommended Dilution:WB 1:500 - 1:2000

Calculated MW:46kDa

Observed MW:Refer to figures

Immunogen:

A synthetic Peptide of human MNDA

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

PYHIN3;

Catalog #:A3963

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:4332

Isotype:IgG

Swiss Prot:P41218

Purity:Affinity purification

For research use only.

Background:

The myeloid cell nuclear differentiation antigen (MNDA) is detected only in nuclei of cells of the granulocyte-monocyte lineage. A 200-amino acid region of human MNDA is strikingly similar to a region in the proteins encoded by a family of interferon-inducible mouse genes, designated lfi-201, lfi-202, and lfi-203, that are not regulated in a cell- or tissue-specific fashion. The 1.8-kb MNDA mRNA, which contains an interferon-stimulated response element in the 5-prime untranslated region, was significantly upregulated in human monocytes exposed to interferon alpha. MNDA is located within 2,200 kb of FCER1A, APCS, CRP, and SPTA1. In its pattern of expression and/or regulation, MNDA resembles IFI16, suggesting that these genes participate in blood cell-specific responses to interferons.

To place an order, please [Click HERE](#).