

## SGCG

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IP

**Recommended Dilution:**WB 1:200 - 1:2000 IHC 1:20 - 1:200 IP 1:20 - 1:50

**Calculated MW:**32kDa

**Observed MW:**Refer to figures

**Immunogen:**

A synthetic Peptide of human SGCG

**Storage Buffer:**

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

A4; MAM; DMDA; SCG3; TYPE; DAGA4; DMDA1; LGMD2C; SCARMD2;

**Catalog #:**A4110

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**6445

**Isotype:**IgG

**Swiss Prot:**Q13326

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes gamma-sarcoglycan, one of several sarcolemmal transmembrane glycoproteins that interact with dystrophin. The dystrophin-glycoprotein complex (DGC) spans the sarcolemma and is comprised of dystrophin, syntrophin, alpha- and beta-dystroglycans and sarcoglycans. The DGC provides a structural link between the subsarcolemmal cytoskeleton and the extracellular matrix of muscle cells. Defects in the encoded protein can lead to early onset autosomal recessive muscular dystrophy, in particular limb-girdle muscular dystrophy, type 2C (LGMD2C).

*To place an order, please [Click HERE](#).*