

## STAM

**Reactivity:** Human Mouse Rat

**Tested applications:** WB

**Recommended Dilution:** WB 1:200 - 1:1000

**Calculated MW:** 59kDa

**Observed MW:** Refer to figures

**Immunogen:**

Recombinant protein of human STAM

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

STAM1; STAM-1;

**Catalog #:** A4198

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 8027

**Isotype:** IgG

**Swiss Prot:** Q92783

**Purity:** Affinity purification

For research use only.

**Background:**

This gene encodes a member of the signal-transducing adaptor molecule family. These proteins mediate downstream signaling of cytokine receptors and also play a role in ER to Golgi trafficking by interacting with the coat protein II complex. The encoded protein also associates with hepatocyte growth factor-regulated substrate to form the endosomal sorting complex required for transport-0 (ESCRT-0), which sorts ubiquitinated membrane proteins to the ESCRT-1 complex for lysosomal degradation. Alternatively spliced transcript variants have been observed for this gene.

*To place an order, please [Click HERE](#).*