

## GALNT4

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:20 - 1:200

**Calculated MW:**67kDa

**Observed MW:**Refer to figures

**Immunogen:**

A synthetic Peptide of human GALNT4

**Storage Buffer:**

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

GALNACT4; GALNAC-T4;

**Catalog #:**A4243

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**8693

**Isotype:**IgG

**Swiss Prot:**Q8N4A0

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes a member of the UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase (GalNAc-T) family of enzymes. GalNAc-Ts initiate mucin-type O-linked glycosylation in the Golgi apparatus by catalyzing the transfer of GalNAc to serine and threonine residues on target proteins. They are characterized by an N-terminal transmembrane domain, a stem region, a luminal catalytic domain containing a GT1 motif and Gal/GalNAc transferase motif, and a C-terminal ricin/lectin-like domain. GalNAc-Ts have different, but overlapping, substrate specificities and patterns of expression. In vitro, the encoded protein can complement other GalNAc-Ts in the complete O-glycosylation of the mucin-1 tandem repeat and can O-glycosylate the P-selectin glycoprotein ligand-1 molecule. The coding region of this gene is contained within a single exon. Fusion transcripts, which combine part of this gene with the 5' exons of the neighboring POC1B (POC1 centriolar protein homolog B) gene, also exist.

*To place an order, please [Click HERE](#).*