

## MonoMethyl-Histone H3-K14

**Reactivity:**Human Mouse Rat Other (Wide Range)

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:20 - 1:50

**Calculated MW:**15kDa

**Observed MW:**Refer to Figures

**Immunogen:**

A synthetic peptide of human MonoMethyl-Histone H3-K14

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

q

**Synonym:**

H3K14me1; H3t; H3.4; H3/g; H3FT;

**Catalog #:**A5277

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**8290

**Isotype:**IgG

**Swiss Prot:**Q16695

**Purity:**Affinity purification

For research use only.

**Background:**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

*To place an order, please [Click HERE](#).*