

## PCSK5

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**207kDa

**Observed MW:**Refer to figures

**Immunogen:**

Recombinant protein of human PCSK5

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

PC5; PC6; PC6A; SPC6;

**Catalog #:**A5450

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**5125

**Isotype:**IgG

**Swiss Prot:**Q92824

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an initial autocatalytic processing event in the ER to generate a heterodimer which exits the ER. It then sorts to the trans-Golgi network where a second autocatalytic event takes place and the catalytic activity is acquired. This encoded protein is widely expressed and one of the seven basic amino acid-specific members which cleave their substrates at single or paired basic residues. It mediates posttranslational endoproteolytic processing for several integrin alpha subunits and is thought to process prorenin, pro-membrane type-1 matrix metalloproteinase and HIV-1 glycoprotein gp160. Alternative splicing results in multiple transcript variants, some of which encode distinct isoforms, including a protease packaged into dense core granules (PC5A) and a type 1 membrane bound protease (PC5B).

**To place an order, please [Click HERE](#).**