

## LITAF

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:**17kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human LITAF

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

b

**Synonym:**

PIG7; SIMPLE; TP53I7;

**Catalog #:**A5469

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**9516

**Isotype:**IgG

**Swiss Prot:**Q99732

**Purity:**Affinity purification

For research use only.

**Background:**

Lipopolysaccharide is a potent stimulator of monocytes and macrophages, causing secretion of tumor necrosis factor-alpha (TNF-alpha) and other inflammatory mediators. This gene encodes lipopolysaccharide-induced TNF-alpha factor, which is a DNA-binding protein and can mediate the TNF-alpha expression by direct binding to the promoter region of the TNF-alpha gene. The transcription of this gene is induced by tumor suppressor p53 and has been implicated in the p53-induced apoptotic pathway. Mutations in this gene cause Charcot-Marie-Tooth disease type 1C (CMT1C) and may be involved in the carcinogenesis of extramammary Paget's disease (EMPD). Multiple alternatively spliced transcript variants have been found for this gene.

*To place an order, please [Click HERE](#).*