

## UBE2C

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:**19kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human UBE2C

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

bfl

**Synonym:**

UBCH10; dJ447F3.2;

**Catalog #:**A5499

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**11065

**Isotype:**IgG

**Swiss Prot:**O00762

**Purity:**Affinity purification

For research use only.

**Background:**

The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, ubiquitin-conjugating enzymes, and ubiquitin-protein ligases. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is required for the destruction of mitotic cyclins and for cell cycle progression, and may be involved in cancer progression. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been defined on chromosomes 4, 14, 15, 18, and 19.

*To place an order, please [Click HERE](#).*