

## UGT1A4

**Reactivity:**Human

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**60kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human UGT1A4

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

UDPGT; UGT1D; UGT-1D; UGT1.4; HUG-BR2; UGT1-04; UDPGT 1-4;

**Catalog #:**A5549

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**54657

**Isotype:**IgG

**Swiss Prot:**P22310

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes a UDP-glucuronosyltransferase, an enzyme of the glucuronidation pathway that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into water-soluble, excretable metabolites. This gene is part of a complex locus that encodes several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons followed by four common exons. Four of the alternate first exons are considered pseudogenes. Each of the remaining nine 5' exons may be spliced to the four common exons, resulting in nine proteins with different N-termini and identical C-termini. Each first exon encodes the substrate binding site, and is regulated by its own promoter. This enzyme has some glucuronidase activity towards bilirubin, although is more active on amines, steroids, and sapogenins.

**To place an order, please [Click HERE](#).**