

## LAT

---

**Reactivity:**Human

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**27kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human LAT

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

LAT1; pp36;

**Catalog #:**A5650

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**27040

**Isotype:**IgG

**Swiss Prot:**O43561

**Purity:**Affinity purification

For research use only.

**Background:**

The protein encoded by this gene is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative splicing results in multiple transcript variants encoding different isoforms.

*To place an order, please [Click HERE](#).*