

NLRP3

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW: 118kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human NLRP3

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

b

Synonym:

FCU; MWS; FCAS; Cias1; Mmig1; NALP3; Pypaf1; Ail/AVP; AGTAVPRL;

Catalog #: A5652

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 114548

Isotype: IgG

Swiss Prot: Q96P20

Purity: Affinity purification

For research use only.

Background:

This gene encodes a pyrin-like protein containing a pyrin domain, a nucleotide-binding site (NBS) domain, and a leucine-rich repeat (LRR) motif. This protein interacts with the apoptosis-associated speck-like protein PYCARD/ASC, which contains a caspase recruitment domain, and is a member of the NALP3 inflammasome complex. This complex functions as an upstream activator of NF-kappaB signaling, and it plays a role in the regulation of inflammation, the immune response, and apoptosis. Mutations in this gene are associated with familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome, and neonatal-onset multisystem inflammatory disease (NOMID). Multiple alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. Alternative 5' UTR structures are suggested by available data; however, insufficient evidence is available to determine if all of the represented 5' UTR splice patterns are biologically valid.

To place an order, please [Click HERE](#).