

## ABCG2

**Reactivity:**Mouse

**Tested applications:**WB

**Recommended Dilution:**WB 1:500 - 1:2000

**Calculated MW:**72kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human ABCG2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

MRX; MXR; ABCP; BCRP; BMDP; MXR1; ABC15; BCRP1; CD338; GOUT1; MXR-1; CDw338; UAQTL1; EST157481;

**Catalog #:**A5661

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**9429

**Isotype:**IgG

**Swiss Prot:**Q9UNQ0

**Purity:**Affinity purification

For research use only.

**Background:**

The membrane-associated protein encoded by this gene is included in the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. Alternatively referred to as a breast cancer resistance protein, this protein functions as a xenobiotic transporter which may play a major role in multi-drug resistance. It likely serves as a cellular defense mechanism in response to mitoxantrone and anthracycline exposure. Significant expression of this protein has been observed in the placenta, which may suggest a potential role for this molecule in placenta tissue. Multiple transcript variants encoding different isoforms have been found for this gene.

*To place an order, please [Click HERE](#).*