

MME

Reactivity: Human Mouse Rat

Tested applications: WB IHC IP FC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IP 1:20 - 1:50 FC 1:20 - 1:50

Calculated MW: 85kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human MME

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

i

Synonym:

NEP; SFE; CD10; CALLA;

Catalog #: A5664

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 4311

Isotype: IgG

Swiss Prot: P08473

Purity: Affinity purification

For research use only.

Background:

This gene encodes a common acute lymphocytic leukemia antigen that is an important cell surface marker in the diagnosis of human acute lymphocytic leukemia (ALL). This protein is present on leukemic cells of pre-B phenotype, which represent 85% of cases of ALL. This protein is not restricted to leukemic cells, however, and is found on a variety of normal tissues. It is a glycoprotein that is particularly abundant in kidney, where it is present on the brush border of proximal tubules and on glomerular epithelium. The protein is a neutral endopeptidase that cleaves peptides at the amino side of hydrophobic residues and inactivates several peptide hormones including glucagon, enkephalins, substance P, neurotensin, oxytocin, and bradykinin. This gene, which encodes a 100-kD type II transmembrane glycoprotein, exists in a single copy of greater than 45 kb. The 5' untranslated region of this gene is alternatively spliced, resulting in four separate mRNA transcripts. The coding region is not affected by alternative splicing.

To place an order, please [Click HERE](#).