

GRN

Reactivity: Human Mouse

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW: 63kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human GRN

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

j

Synonym:

GEP; GP88; PEPI; PGRN; CLN11; PCDGF;

Catalog #: A5773

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 2896

Isotype: IgG

Swiss Prot: P28799

Purity: Affinity purification

For research use only.

Background:

Granulins are a family of secreted, glycosylated peptides that are cleaved from a single precursor protein with 7.5 repeats of a highly conserved 12-cysteine granulin/epithelin motif. The 88 kDa precursor protein, progranulin, is also called proepithelin and PC cell-derived growth factor. Cleavage of the signal peptide produces mature granulin which can be further cleaved into a variety of active, 6 kDa peptides. These smaller cleavage products are named granulin A, granulin B, granulin C, etc. Epithelins 1 and 2 are synonymous with granulins A and B, respectively. Both the peptides and intact granulin protein regulate cell growth. However, different members of the granulin protein family may act as inhibitors, stimulators, or have dual actions on cell growth. Granulin family members are important in normal development, wound healing, and tumorigenesis.

To place an order, please [Click HERE](#).