

## APEH

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

**Calculated MW:**81kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human APEH

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

j

**Synonym:**

APH; OPH; AARE; ACPH; D3S48E; D3F15S2; DNF15S2;

**Catalog #:**A5893

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**327

**Isotype:**IgG

**Swiss Prot:**P13798

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes the enzyme acylpeptide hydrolase, which catalyzes the hydrolysis of the terminal acetylated amino acid preferentially from small acetylated peptides. The acetyl amino acid formed by this hydrolase is further processed to acetate and a free amino acid by an aminoacylase. This gene is located within the same region of chromosome 3 (3p21) as the aminoacylase gene, and deletions at this locus are also associated with a decrease in aminoacylase activity. The acylpeptide hydrolase is a homotetrameric protein of 300 kDa with each subunit consisting of 732 amino acid residues. It can play an important role in destroying oxidatively damaged proteins in living cells. Deletions of this gene locus are found in various types of carcinomas, including small cell lung carcinoma and renal cell carcinoma.

**To place an order, please [Click HERE](#).**