

RPL10A

Reactivity:Human Mouse Rat

Tested applications:WB

Recommended Dilution:WB 1:1000 - 1:2000

Calculated MW:25kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human RPL10A

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

j

Synonym:

L10A; CSA19; NEDD6; Csa-19;

Catalog #:A5925

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:4736

Isotype:IgG

Swiss Prot:P62906

Purity:Affinity purification

For research use only.

Background:

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L1P family of ribosomal proteins. It is located in the cytoplasm. The expression of this gene is downregulated in the thymus by cyclosporin-A (CsA), an immunosuppressive drug. Studies in mice have shown that the expression of the ribosomal protein L10a gene is downregulated in neural precursor cells during development. This gene previously was referred to as NEDD6 (neural precursor cell expressed, developmentally downregulated 6), but it has been renamed RPL10A (ribosomal protein 10a). As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

To place an order, please [Click HERE](#).