

DDX17

Reactivity:Human

Tested applications:WB IHC FC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100 FC1:20 - 1:50

Calculated MW:80kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human DDX17

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

j

Synonym:

P72; RH70

Catalog #:A5957

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:10521

Isotype:IgG

Swiss Prot:Q92841

Purity:Affinity purification

For research use only.

Background:

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure, such as translation initiation, nuclear and mitochondrial splicing, and ribosome and splicesosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein, which is an ATPase activated by a variety of RNA species, but not by dsDNA. This protein, and that encoded by DDX5 gene, are more closely related to each other than to any other member of the DEAD box family. This gene can encode multiple isoforms due to both alternative splicing and the use of alternative translation initiation codons, including a non-AUG (CUG) start codon.

To place an order, please [Click HERE](#).