

## METAP2

**Reactivity:** Human Mouse

**Tested applications:** WB

**Recommended Dilution:** WB 1:200 - 1:2000

**Calculated MW:** 53kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human METAP2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

p67; MAP2; MNPEP; p67eIF2;

**Catalog #:** A5962

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 10988

**Isotype:** IgG

**Swiss Prot:** P50579

**Purity:** Affinity purification

For research use only.

**Background:**

This gene is a member of the methionyl aminopeptidase family and encodes a protein that binds 2 cobalt or manganese ions. This protein functions both by protecting the alpha subunit of eukaryotic initiation factor 2 from inhibitory phosphorylation and by removing the amino-terminal methionine residue from nascent protein. Increased expression of this gene is associated with various forms of cancer and the anti-cancer drugs fumagillin and ovalicin inhibit the protein by irreversibly binding to its active site. A pseudogene of this gene is located on chromosome 2.

**To place an order, please [Click HERE](#).**