

CTCFL

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:20 - 1:100

Calculated MW:75kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human CTCFL

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

CT27; BORIS; CTCF-T; HMGB1L1; dJ579F20.2;

Catalog #:A6149

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:140690

Isotype:IgG

Swiss Prot:Q8NI51

Purity:Affinity purification

For research use only.

Background:

CCCTC-binding factor (CTCF), an 11-zinc-finger factor involved in gene regulation, utilizes different zinc fingers to bind varying DNA target sites. CTCF forms methylation-sensitive insulators that regulate X-chromosome inactivation. This gene is a paralog of CTCF and appears to be expressed primarily in the cytoplasm of spermatocytes, unlike CTCF which is expressed primarily in the nucleus of somatic cells. CTCF and the protein encoded by this gene are normally expressed in a mutually exclusive pattern that correlates with resetting of methylation marks during male germ cell differentiation. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene.

To place an order, please [Click HERE](#).