

## MSLN

**Reactivity:**Human

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

**Calculated MW:**68kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human MSLN

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

MPF; SMRP;

**Catalog #:**A6183

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**10232

**Isotype:**IgG

**Swiss Prot:**Q13421

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes a precursor protein that is cleaved into two products, megakaryocyte potentiating factor and mesothelin. Megakaryocyte potentiation factor functions as a cytokine that can stimulate colony formation in bone marrow megakaryocytes. Mesothelin is a glycosylphosphatidylinositol-anchored cell-surface protein that may function as a cell adhesion protein. This protein is overexpressed in epithelial mesotheliomas, ovarian cancers and in specific squamous cell carcinomas. Alternative splicing results in multiple transcript variants.

*To place an order, please [Click HERE](#).*