

## COMT

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC IF

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:** 30kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human COMT

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

1 mg/ml

**Synonym:**

HEL-S-98n;

**Catalog #:** A6200

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 1312

**Isotype:** IgG

**Swiss Prot:** P21964

**Purity:** Affinity purification

For research use only.

**Background:**

Catechol-O-methyltransferase catalyzes the transfer of a methyl group from S-adenosylmethionine to catecholamines, including the neurotransmitters dopamine, epinephrine, and norepinephrine.

This O-methylation results in one of the major degradative pathways of the catecholamine transmitters. In addition to its role in the metabolism of endogenous substances, COMT is important in the metabolism of catechol drugs used in the treatment of hypertension, asthma, and Parkinson disease. COMT is found in two forms in tissues, a soluble form (S-COMT) and a membrane-bound form (MB-COMT). The differences between S-COMT and MB-COMT reside within the N-termini. Several transcript variants are formed through the use of alternative translation initiation sites and promoters.

**To place an order, please [Click HERE](#).**