

## CYP2C9

**Reactivity:**Human

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

**Calculated MW:**55kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human CYP2C9

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

CPC9; CYP2C; CYP2C10; CYP11C9; P45011C9;

**Background:**

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by rifampin. The enzyme is known to metabolize many xenobiotics, including phenytoin, tolbutamide, ibuprofen and S-warfarin. Studies identifying individuals who are poor metabolizers of phenytoin and tolbutamide suggest that this gene is polymorphic. The gene is located within a cluster of cytochrome P450 genes on chromosome 10q24.

*To place an order, please [Click HERE](#).*

**Catalog #:**A6219

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**1559

**Isotype:**IgG

**Swiss Prot:**P11712

**Purity:**Affinity purification

For research use only.